




Abortion care:

Information and legal
obligations for medical
practitioners

One team, **many** dreams.

Care / Respect / Innovation / Teamwork / Integrity



Contents

Introduction	4
Abortion and the law in Western Australia	5
Abortion before 20 weeks	6
Abortion from 20 weeks	6
Requirements of informed consent	7
Capacity to consent	8
Requirements for dependant minors	9
When is a young person not considered a dependant minor?	9
Dependant minors and parental involvement	9
Obtaining a Children's Court Order	10
How is a Children's Court order obtained?	10
How is the medical practitioner involved in the Children's Court order?	11
Dependant who is unable to give informed consent	11
Pregnancy following sexual assault	12
Self-managed abortion with unregulated medication	13
Requirement to notify the Chief Health Officer	13
Ethical obligations	14
Understanding your obligations	14
Quick reference guide	16
Abortion and pregnancy information	17
Methods of induced abortion	17
Medical abortion	17
Surgical abortion	17
Availability of medical and surgical abortion	18

Risks of induced abortion	19
Short-term risks and complications of medical and surgical abortion	19
Long-term complications	22
Risks of carrying a pregnancy to term	24
Mortality risk	24
Morbidity	24
Other risks associated with pregnancy	25
Table 1: Overview of medical risks of abortion and continuing the pregnancy	26
Adoption	27
Guidelines for medical practitioner counselling	28
Resources	29
Abortion providers	29
Unplanned pregnancy counselling	30
Legal services	30
Womens health services	31
Mental health services	31
Other contacts	32
Appendix A: Suggested further information points for counselling on the processes involved	33
Appendix B: Abortion referral process template	34
Notification by medical practitioner of induced abortion	35
References	36
Disclaimer	39
Copyright	39

Introduction

This booklet aims to:

- Increase awareness of the requirements for an abortion in Western Australia;
- Provide an overview of medical risks of both abortion and pregnancy; and
- List options for accessing pregnancy and abortion support services.

Medical practitioners should be aware that:

- Abortion is a safe procedure
- Abortion is legal prior to 20 weeks gestation in Western Australia, providing conditions of informed consent are met
- Abortion is only legal from 20 weeks gestation in Western Australia in specific circumstances; and
- There are two options available for abortion, medical and surgical.

Medical practitioners should also consider:

- Arranging a dating ultrasound scan as soon as possible if there is uncertainty about the date of first day of last normal menstrual period
- Providing an appropriate environment for assessing the pregnant women to ensure as far as possible that no coercion or pressure has been applied; and
- That there is a cost associated with abortion. In situations of financial distress, financial support may be available by referral to the Family Services Coordinator at King Edward Memorial Hospital (KEMH) KEMH.Referrals@health.wa.gov.au or fax (08) 6458 1031. KEMH will contact the patient via telephone and conduct a needs assessment.



Abortion and the law in Western Australia

In May 1998, amendments to legislation, including the Criminal Code 1913 and the Health Act 1911, known from 2016 onwards as the *Health (Miscellaneous Provisions) Act 1911* (WA), enabled the lawful performance of an abortion when:

- The abortion is performed by a medical practitioner in good faith and with reasonable care and skill; and
- The performance of the abortion is justified under section 334 of the *Health (Miscellaneous Provisions) Act 1911* (WA).

Under section 334 of the *Health (Miscellaneous Provisions) Act 1911* (WA), the performance of an abortion is justified for the purposes of section 199(1) of the Criminal Code if:

- a) the woman concerned has given informed consent; or
- b) the woman concerned will suffer serious personal, family or social consequences if the abortion is not performed; or
- c) serious danger to the physical or mental health of the woman concerned will result if the abortion is not performed; or
- d) the pregnancy of the woman concerned is causing serious danger to her physical or mental health.



Abortion before 20 weeks

Abortion is available on request before 20 weeks of pregnancy provided that informed consent has been given.

There is always a balance between referral early in pregnancy and allowing sufficient time for decision-making. However, it is important to ensure that women wanting abortion care are referred early, as the risk of complications from the procedure, and costs, rise with increasing gestation.

Abortion from 20 weeks

Section 334(7) *Health (Miscellaneous Provisions) Act 1911* (WA) states that:

If at least 20 weeks of the woman's pregnancy have been completed when the abortion is performed, the performance of the abortion is not justified unless:

a) Two medical practitioners who are members of a panel of at least six medical practitioners, by appointment by the Minister for the purposes of this section have agreed that the mother, or the unborn child, has a severe medical condition that in the clinical judgment of those two medical practitioners, justifies the procedure; and

b) The abortion is performed in a facility approved by the Minister for the purposes of this section.

If a woman is requesting an abortion which may need to occur from 20 weeks gestation the referring medical practitioner must contact the King Edward Memorial Hospital family services co-ordinator on (08)6458 2222. The co-ordinator will discuss how to proceed with an application to the panel and facilitate the necessary Maternal-Fetal Medicine assessment or perinatal psychiatry assessment. Note:

- The referring medical practitioner is still required to obtain the woman's informed consent.
- Abortion from 20 weeks gestation can only be performed at a facility approved by the Minister for Health; these are King Edward Memorial Hospital and Broome Regional Hospital.

Requirements of informed consent

Western Australia's abortion legislation sets out the "informed consent" that a woman must freely give prior to an abortion. A medical practitioner responsible for this "informed consent" cannot also perform or assist with the abortion. This effectively results in two separate requirements, where, after discussion with the first medical practitioner, the woman wishes to proceed with the procedure.

1. The *Health (Miscellaneous Provisions) Act 1911* (WA) defines informed consent, for this purpose, as consent freely given by the woman where a medical practitioner has:
 - properly, appropriately and adequately provided her with counselling about the medical risk of abortion and of carrying a pregnancy to term; and
 - offered her the opportunity of referral to appropriate and adequate counselling about matters relating to abortion and carrying a pregnancy to term; and
 - informed her that appropriate and adequate counselling will be available to her should she wish it, following an abortion or after carrying the pregnancy to term.

If the woman does not provide informed consent as above, no referral should be made.

Medical practitioners should note that the term 'counselling' in this case is synonymous with providing information; it is not psychological counselling to assist with decision-making about pregnancy choices. Although many doctors would see supportive counselling as part of their role, it is not a legal requirement in relation to informed consent.

- This booklet provides information to assist in properly, appropriately and adequately providing her with counselling on risks related to pregnancy and abortion; see 'Abortion and Pregnancy information' below.
 - Information is also provided on organisations funded to provide unintended pregnancy counselling at no cost to the woman, should she decide to take up the offer of referral to counselling. See a list of providers in the Resource section at the back of this booklet under 'Unplanned pregnancy counselling'.
 - Many medical practitioners may provide counselling themselves but are also obliged to offer the opportunity of referral. Whether or not such an offer is taken up is a matter for the woman, i.e. she does not have to be counselled elsewhere in order to meet the requirements. A brief guide outlining the principles of counselling can be found on page 28 of this booklet.
2. The second medical practitioner must not perform an abortion unless they are satisfied that the woman has given informed consent (as understood at common law) to the procedure. That is, consent must be voluntary (the decision must be made without duress or coercion); the person giving it must have capacity to give it; it must be informed (with discussion of maternal risks); it must cover the procedure to be performed, and it must be current. A woman may withdraw her consent at any time prior to the procedure being performed. More detail can be found at the Consent to Treatment Policy.

Capacity to consent

Section 334(4) of the *Health (Miscellaneous Provisions) Act 1911* (WA) provides that where it is impracticable for a woman to give informed consent, the performance of an abortion will be justified (without such consent) if, and only if:

- serious danger to the physical or mental health of the woman concerned will result if an abortion is not performed; or
- the pregnancy of the woman concerned is causing serious danger to her physical or mental health.

The informed consent required under s.334 *Health (Miscellaneous Provisions) Act 1911* (WA) can only be given by 'the woman concerned', unless it is impracticable as stated above. This means that a guardian, even where formally appointed under the Guardianship and Administration Act 1990 (WA) to make treatment decisions on behalf of a woman, cannot provide the required consent on behalf of the woman (see the decision of the State Administrative Tribunal (SAT) - KS and CL 2015 WASAT 9). Note, however, that a person's capacity can vary depending on the proposed treatment. Depending on the facts of a case, a woman may not have capacity in respect of some treatment decisions, but the SAT, in examining all the facts of the case, may determine she has sufficient capacity to provide informed consent in relation to a proposed abortion.

Accordingly, if a medical practitioner is faced with a situation where an abortion is proposed for a woman who potentially lacks decision-making capacity, it is recommended that the medical practitioner seeks urgent legal advice or, in the case of adults, makes an urgent application for a hearing at SAT (for SAT to review the decision-making capacity of 'the woman concerned').

Dependant minors

Section 334(8) of the *Health (Miscellaneous Provisions) Act 1911* (WA) states:

8. For the purposes of this section —

- a) subject to subsection (11), a woman who is a dependant minor shall not be regarded as having given informed consent unless a custodial parent of the woman has been informed that the performance of an abortion is being considered and has been given the opportunity to participate in a counselling process and in consultations between the woman and her medical practitioner as to whether the abortion is to be performed;
- b) a woman is a dependant minor if she has not reached the age of 16 years and is being supported by a custodial parent or parents; and
- c) a reference to a parent includes a reference to a legal guardian.

When is a young person not considered a dependant minor?

If the young person is under 16 years and is not being supported by a custodial parent, the requirements for dependant minors outlined above do not apply.

The legislation does not define what is meant by “supported”. However, it could be reasonable to interpret it as referring primarily to financial support and a child living away from home who is not financially dependent on her parents would not be a ‘dependant minor’.

When minors are considered independent, the WA legislation on abortion is the same as for people over 16 years of age.

Dependant minors and parental involvement

It should be noted that the legal requirement is only that a custodial parent is given the opportunity to participate in counselling/consultation. Whether or not this opportunity is taken is a matter for them. The medical practitioner should be satisfied that the custodial parent has been informed and invited to become involved in counselling and consultations. Alternatively, a dependant minor may make an application to the Children’s Court to waive this requirement, see below.

Medical practitioners should note that it is only in informing the custodial parent and giving them the opportunity to participate in consultation that an exception to normal patient confidentiality exists. In all other aspects related to the abortion and care the usual requirements of medical practitioner/patient confidentiality apply (i.e. that confidentiality is maintained by the medical practitioner except where the dependant minor has consented to the release of information).

Free legal assistance for those who decide to pursue this option is available from the Children's Court Protection Service. Their contact details are in the Resources lists at the end of this booklet. This service will assist the woman with applications to the Children's Court, including helping with the completion of a form available from the Court and accompanying the woman to Court to put her case to the Magistrate.

How is a Children's Court order obtained?

A woman who is a dependant minor may apply to the Children's Court for an order that a person specified in the application, being a custodial parent of the woman, should not be given the information and opportunity referred to in subsection (8)(a) and the court may, on being satisfied that the application should be granted, make an order in those terms.

The decision as to whether to inform the custodial parent, or to seek to vary this requirement by applying to the Children's Court under section 334(9) of the *Health (Miscellaneous Provisions) Act 1911* (WA) is one for the dependant minor herself to make. The requirements of law in relation to dependant minors may be varied by an order of the Children's Court under Section 334(9) of the *Health (Miscellaneous Provisions) Act 1911* (WA) which states:

Obtaining a Children's Court Order

- Where a medical practitioner considers that a patient may be under the age of 16 years, it is strongly recommended that the medical practitioner seek some proof of age.
- Where the young person is under the age of 16 years, it will also be necessary for a medical practitioner to determine whether or not they are being supported by a custodial parent.
- If the custodial parent is provided with the necessary information and has been given the opportunity to participate in counselling and consultations the decision to proceed with referral is the decision of the young person.
- It is possible that a dependant minor may be able to give the necessary informed consent, even if this is not consistent with the custodial parent's views.

Some points to consider:

How is the medical practitioner involved in the Children's Court order?

Medical practitioners can be guided by a legal service, which the young person may consult, as to the information and actions required in relation to the court process. In such cases, the medical practitioner will usually be asked to provide a letter which contains an assessment of the maturity of the young person and her social circumstances in so far as they may be known to the medical practitioner. Such a letter would generally be provided by the legal service to the magistrate for the purpose of assisting the magistrate to make a decision on the application. The application is generally heard within a few days by a Magistrate and a decision made.

If the magistrate makes an order that a custodial parent should not be given the information and opportunity referred to in section 334(8)(a) of the *Health (Miscellaneous Provisions) Act 1911(WA)*, then informed consent can be given by the woman as long as the usual requirements of section 334(5) of the *Health (Miscellaneous Provisions) Act 191 (WA)* have been met.

Medical practitioners should note that in this situation extra support may be required, especially where there is little family support. However, it is also important that medical practitioners keep in mind that any decision to apply to the Children's Court is ultimately one for the young person to make, not the medical practitioner.

Dependant who is unable to give informed consent

Section 334(4) of the *Health (Miscellaneous Provisions) Act 1911 (WA)* provide that where it is impracticable for a woman to give informed consent, the performance of an abortion will be justified without such consent where:

- serious danger to the physical or mental health of the woman will result if an abortion is not performed; or
- the pregnancy of the woman is causing serious danger to her physical or mental health.

If a medical practitioner is concerned about the capacity of a woman to give consent for referral, it may be appropriate to apply to the Children's Court for permission for the abortion to be carried out. A referral to a legal service may be required in such situations.

Pregnancy following sexual assault

Unwanted pregnancy may be the result of a recent sexual assault. If there is disclosure of a sexual assault it is important to listen to and believe the victim.

Points to consider:

- Determine accurate gestation. An ultrasound will confirm gestation and allow correlation with alleged date of incident.

- Ask the patient if they would like a referral to sexual assault counselling. This can be done through the Sexual Assault Resource Centre (SARC), Family Services or Sexual Health Quarters.

- Contact SARC duty officer on 08 6458 1820 if you would like more individualised advice.
- Ask the patient if they would like police involvement. Note that products of conception can be used as DNA evidence and this can be requested by police and taken as evidence.

- It is possible that the patient is at ongoing risk of harm. Follow the Shared Maternity Care provider (WA) - Referral Pathway for Family and Domestic Violence (page 12-13) to screen for FDV and complete the appropriate risk assessment.

- For more information on sexual assault, see the Sexual Assault Resource Centre website which has information for clients and health professionals.

Under 18 years of age

If the medical practitioner has a reasonable belief that a person under 18 years of age has been sexually abused, a mandatory report to the Child Protection Unit is required, even when the patient is consider a mature minor. See Mandatory reporting of child abuse and neglect | Child Family Community Australia (aifs.gov.au)

Self-managed abortion with unregulated medication

With the increased availability of abortion medicines via the internet it is possible medical practitioners will see women who have attempted, or intend to attempt, self-managed abortion without clinical supervision ⁽¹⁾.

It is important that women are aware of the need for medical supervision and appropriate medications for safe and effective abortion. Some online abortion medications are unregulated and may be counterfeit ⁽²⁾. Risks include failed treatment, health risks to the woman and risk to subsequent pregnancies ⁽³⁾. If a woman has used unregulated medication for a self-managed abortion, she should be encouraged to seek medical treatment.

Requirement to notify the Chief Health Officer

Any medical practitioner who performs an abortion must notify the Chief Health Officer of the event within 14 days of the abortion being performed. This includes medical practitioners who provide medical abortions or surgical abortions in community settings. Such notifications are not required by the medical practitioner making the referral for abortion.

Notification must be made using Form 1 – Notification by Medical Practitioner of Induced Abortion. <https://datalibrary-rc.health.wa.gov.au/surveys/?s=CAR9J78MRT>

If a paper version of the form is preferred, or for more information required, contact Maternal and Child Health, Data Management, Information and Performance Governance Unit on 9222 2417 or birthdata@health.wa.gov.au

For more information, see https://ww2.health.wa.gov.au/Articles/A_E/Abortion-Notification-System

Ethical obligations

Understanding your obligations

Medical practitioners are not required by legislation to participate in consultation and referral for abortion. However, they should be aware of their obligations as outlined in the Australian Health Practitioner Regulation Agency (AHPRA) and Medical Board of Australia's Good Medical Practice: a code of conduct for doctors in Australia.

The relevant sections of the Good medical practice: a code of conduct for doctors in Australia 2020 include:

3.4 Decisions about access to medical care Your decisions about patients' access to medical care must be free from bias and discrimination. Good medical practice involves:

3.4.1 Treating your patients with respect at all times.

3.4.2 Not prejudicing your patient's care because you believe that a patient's behaviour has contributed to their condition.

3.4.3 Upholding your duty to your patient and not discriminating against your patient on grounds such as race, religion, sex, gender identity, sexual orientation, disability or other grounds, as described in antidiscrimination legislation.

3.4.6 Being aware of your right to not provide or directly participate in treatments to which you conscientiously object, informing your patients and, if relevant, colleagues of your objection, and not using your objection to impede access to treatments that are legal. In some jurisdictions, legislation mandates doctors who do not wish to participate in certain treatments, to refer on the patient.

3.4.7 Not allowing your moral or religious views to deny patients access to medical care, recognising that you are free to decline to personally provide or directly participate in that care.

Good medical practice: a code of conduct for doctors in Australia complements the Australian Medical Association Code of Ethics. The relevant section states:


2.1.13 If you refuse to provide or participate in some form of diagnosis or treatment based on a conscientious objection, inform the patient so that they may seek care elsewhere. Do not use your conscientious objection to impede patients' access to medical treatments, including in an emergency situation.

The AMA Position Statement on conscientious objection 2019 states:

"2.3 A doctor with a conscientious objection, should:

- inform the patient of their objection, preferably in advance or as soon as practicable;
- inform the patient that they have the right to see another doctor and ensure the patient has sufficient information to enable them to exercise that right;
- take whatever steps are necessary to ensure the patient's access to care is not impeded;
- continue to treat the patient with dignity and respect, even if the doctor objects to the treatment or procedure the patient is seeking;
- continue to provide other care to the patient, if they wish;
- refrain from expressing their own personal beliefs to the patient in a way that may cause them distress;
- inform their employer, or prospective employer, of their conscientious objection and discuss with their employer how they can practice in accordance with their beliefs without compromising patient care or placing a burden on their colleagues."

If a medical practitioner cannot provide the services requested, it may be appropriate to refer the client to a suitable service. Such services are listed under 'Unplanned pregnancy counselling' in the Resources section of this booklet.




Gestation

Age of informed consent

Meeting your obligation

Defining your role



Abortion and pregnancy information

Methods of induced abortion

A pregnancy may be ended using surgical or medical techniques or a combination of both.

Medical abortion

Medical abortion refers to the use of medication to terminate a pregnancy.

The registration by the Pharmaceutical Benefits Scheme of mifepristone for use in Australia for early medical abortion has enabled abortion provision up to 63 days of gestation in an outpatient environment ⁽⁴⁾. General practitioners and hospital doctors can only prescribe mifepristone for this purpose after successfully completing an online course provided by the PBS sponsor, Marie Stopes Australia (www.ms2step.com.au).

Medical abortion involves the use of two agents - mifepristone, a synthetic anti progesterone, and misoprostol, a prostaglandin analogue. The procedure involves oral mifepristone which inhibits the action of progesterone in maintaining the pregnancy and therefore cause the embryo and placental sac to separate from the wall of the uterus. Misoprostol, taken 24-48 hours after mifepristone, induces contractions, cervical opening and causes the evacuation of contents of the uterus.

The combination of mifepristone and misoprostol for women in early pregnancy results in complete abortion in 95 percent of cases ⁽⁵⁾. The other 5 percent of women may need surgical evacuation for retained products of conception ⁽⁴⁾.

Medical abortion can be offered in a primary care setting or through a clinic (see Abortion Providers in Resources). After nine weeks gestation medical abortion is not available in the community but may be available as a hospital inpatient ⁽⁴⁾.

Women undergoing medical abortion will have medical supervision and access to surgical treatment if required. Having a further procedure may incur a further cost.

Surgical abortion

Surgical methods include suction curettage (vacuum aspiration) or dilation and evacuation (D&E). Aspiration techniques can generally be used up to 12 to 14 weeks; however, as gestation increases, safe removal requires cervical preparation and a combination of techniques to remove the products from the uterus ⁽⁶⁾. Surgical abortion is available in licensed day surgery clinics and in hospitals.



Availability of medical and surgical abortion

- Suitability for medical or surgical abortion requires consideration of a range of factors including gestation; the woman's individual circumstances (eg: psychological impacts, social support); medical conditions; and the choice of abortion provider. All evidence indicates that both medical and surgical abortion are safe. Possible adverse effects are outlined below.
- Routine abortion is generally provided through private abortion clinics in the community. The provider may offer early medical abortion as an outpatient or a choice of medical or surgical abortion at a clinic. Medical abortion up to nine weeks gestation can also be provided by trained general practitioners⁽⁷⁾.
- Medical abortion can be available in a hospital setting after nine weeks as a safe alternative to a surgical procedure and may avoid the need for anaesthetic. Contact the local public gynaecological service provider or King Edward Memorial Hospital family services co-ordinator.
- King Edward Memorial Hospital does not provide a routine abortion service; however, it does have an abortion service to assist patients who are unsuitable for private abortion clinics due to medical co-morbidities or are requesting a medical abortion after nine weeks' gestation.
- Any young woman under 14 years of age requesting an abortion should be referred to King Edward Memorial Hospital. This is a specialised service available to all young women under 14 years from across WA.
- Women with restricted financial circumstances that would preclude them from accessing private abortion providers may also be referred to King Edward Memorial Hospital for assistance. This will involve a consultation conducted via telephone between the family services co-ordinator and the woman via a referral from their GP.

See the contacts at the end of the booklet for a list of abortion providers.

Risks of induced abortion

The following sections are an evidence-based summary of the literature on the risks of abortion and of continuing the pregnancy to term.

All of the available evidence indicates that induced abortion both via medical or surgical methods, especially in early pregnancy, is a low-risk procedure ⁽⁶⁾. The risks of death and serious complication with induced abortion are lower than the risks of carrying a pregnancy to term ^(8,9). There are many issues for women to consider when deciding to have an abortion. The medical risks of abortion and continuation of pregnancy is one part of this potentially complex decision.

Short-term risks and complications of medical and surgical abortion

- **Mortality risk**

Both medical and surgical abortion are safe procedures. At all gestational ages, major complications and mortality are rare ⁽¹⁰⁾.

The risk of maternal death from an induced abortion performed by a trained clinician is much lower than carrying a pregnancy to term ⁽⁶⁾. However, mortality increases with gestational age, from 0.1 per 100 000 at eight weeks gestation to 8.9 per 100 000 at 21 weeks ⁽⁶⁾.

- **Morbidity risk**

Despite increased risk with gestational age, rates of complication remain low and are comparable between medical and surgical methods of abortion ^(6,10).

- **Haemorrhage**

The risk of haemorrhage following abortion is low ^(10,11). Estimates of haemorrhage following vacuum aspiration in the first trimester range from 0 to 3 per 1000 cases ⁽¹¹⁾. The risk of blood transfusion following a medical abortion is approximately 0.1 percent ⁽¹²⁾. Risk increases with increased gestation for both medical and surgical abortion ⁽⁶⁾.

- **Infection**

Routine prophylactic antibiotics are offered to women who are undergoing surgical abortion as recommended by the World Health Organisation and the Royal College of Obstetrics and Gynaecology (RCOG). Infection occurs in 0 to 2 percent of cases of surgical and less than 1 percent of cases in medical abortion ^(6,10,13).

- **Retained products of conception**

Medical and surgical methods are generally effective in completing the abortion; however, there is a small risk (less than 2 in 100 for surgical and 5 in 100 for medical) of the need for further intervention to complete the procedure ^(14,15).

- **Failure of abortion**

Both medical and surgical abortion carry a small risk of failure to end the pregnancy (1 or 2 in 100), resulting in a further procedure^(14, 16). The risk of failure is higher in very early pregnancy. For terminations performed by suction curettage, there is a three-fold higher failure rate for those performed before seven weeks gestation compared with those performed at seven to 12 weeks gestation⁽¹⁷⁾. There is some research to suggest that failure of medical abortion increased with the woman's age and gestation⁽⁵⁾.

- **Rhesus isoinmunisation**

The administration of Rh (D) immunoglobulin (Anti-D) can reduce the risk of sensitisation and adverse consequences in subsequent pregnancies. Guidelines for the use of Rh (D) immunoglobulin (Anti-D) in obstetrics by the Royal Australian and New Zealand College of Obstetricians and Gynaecologists (RANZCOG) state that all Rh(D) negative women (who have not actively formed their own Anti-D) should be offered Anti-D at medical or surgical abortion within 72 hours of the procedure⁽¹⁸⁾.

There is no international consensus on the use of Anti-D in the first trimester⁽¹⁰⁾. During the COVID-19 pandemic, RANZCOG recognised that testing and administering Anti-D based on the National Institute for Health and Care Excellence (NICE) guideline 2019, potentially adds delay and barriers to care. The college provided a revised statement, to recommend that rhesus status determination and Anti-D are not required for early medical abortion up to 10 weeks⁽¹⁹⁾.

- **Bleeding and cramping (medical abortion only)**

As medical abortion involves the expulsion of products of conception, women should expect to start bleeding within a few hours of administration of the second medication. Some pain and cramping are also to be expected. The amount of bleeding is greater with medical abortion than surgical and appears to increase with gestational age⁽²⁰⁾. Other common adverse effects include nausea, weakness, headache, and dizziness⁽¹²⁾.

- **Effects of prostaglandins (surgical only)**

Cervical priming with prostaglandins reduces the risk of damage to the cervix. Where prostaglandins are used, such as for cervical priming, they can be associated with side effects such as diarrhoea, nausea, vomiting, dizziness, warm flushes, chills or headaches or pain caused by contractions^(21, 22).

- **Complications related to anaesthesia (surgical abortion only)**

A range of anaesthetics, analgesics and techniques can be employed during an abortion, including general anaesthetic, conscious sedation and local anaesthesia. The preferred option depends on gestation, technique, the woman's preferences and the expertise of the service provider.

In Western Australia the most common technique during surgical abortion is conscious sedation, otherwise known as "twilight sedation", which is associated with less post-operative nausea and vomiting⁽²³⁾ and earlier recovery from anaesthesia⁽²⁴⁾.

Conscious sedation is a state of depressed consciousness that allows protective reflexes and the airway to be maintained. Patients can respond appropriately to physical and verbal stimulation and some memory of what has occurred is possible, but it is usually not distressing. Midazolam, fentanyl and propofol are commonly used. Midazolam may temporarily impair the acquisition of new information (anterograde amnesia), while having little effect on previously stored information (retrograde amnesia) ⁽²⁵⁾.

Although less common than when general anaesthesia is used, drowsiness and dizziness can occur after this method ⁽²⁶⁾. Anxiolytics and narcotics used for conscious sedation may cause respiratory depression, especially when they are used together with higher medication doses. There is a risk that the woman may lose her ability to protect her airway ⁽²⁷⁾.

In pregnancies less than 12 weeks gestation the procedure is low risk and usually takes under 15 minutes. The risk of anaesthetic complications is therefore low, but as with all anaesthetics, risk may be increased in the presence of obesity, smoking, diabetes and other chronic illnesses.

- **Injury**

- Uterine perforation (surgical abortion only)

The risk of uterine perforation is low (0.2- 0.8 percent) although it increases with advancing gestation ⁽⁶⁾.

- Cervical trauma (surgical abortion only)

Risk of cervical trauma is linked to gestation age and provider experience and may occur as part of the procedure ⁽⁶⁾. Cervical priming is recommended by organisations such as the National Institute of Health Care and Excellence to prevent injury to the cervix and uterus ^(28, 29).

- **Complications of induced abortion at 12 to 19 weeks**

While second trimester abortions are safe, there is an increase in complications with increasing gestation. Both medical and surgical methods can be used; however, more training is required for surgical abortion at this gestation ⁽³⁰⁾. The main complication of second trimester medical abortion is retained products of conception causing ongoing bleeding ⁽³¹⁾ or necessitating anaesthetic procedure.

Long-term complications

The following section provides a brief review of the evidence relating to long-term complications after an abortion. It focuses on three issues: future reproduction, breast cancer and psychosocial outcomes.

• Effect on future reproduction

The possible long-term adverse effects of abortion on future reproduction can be of particular concern to women. Many women plan to have children in the future. They should be assured that abortion is not associated with an increased risk of infertility⁽³²⁾. The following rare complications can impact adversely upon future fertility: cervical weakening, scarring and stenosis, Asherman's syndrome, post-infection fallopian tube damage, and hysterectomy following post-abortion complications⁽³³⁻³⁵⁾. Induced abortion has been associated with a small increased risk of subsequent preterm birth. This increases with the number of abortions⁽³⁶⁾.

• Breast cancer

Abortion is not associated with an increase in breast cancer⁽³⁶⁾. While there was previously conflicting evidence, it has now been clearly demonstrated that there is no increased risk⁽³⁷⁾.

• Psychological consequences

Women request abortions for multiple and complex reasons related to their individual circumstances that can include socioeconomic status, age, health, parity, marital status, reproductive coercion and intimate partner violence^(38,39). Historically there has been some concern that women who have an abortion will suffer lasting adverse emotional or mental health consequences. However, there is no evidence to suggest this is the case. While it can be expected that women will experience a range of emotions in relation to an abortion, there is no evidence linking abortion to short or long-term negative emotions. Studies indicate that relief is the most common short-term emotion experienced following an abortion; however, all emotions in relation to an abortion decline in intensity with time⁽⁴⁰⁾. Emotions related to abortion are also the product of the individual and social context that the woman is part of, rather than the procedure itself⁽⁴¹⁾. It is important to be aware of the stigma that can be associated with abortion as this can impact on women's wellbeing⁽⁴²⁾.

The American Psychological Association Task Force on Mental Health and Abortion noted that many studies into psychological outcomes of abortion were methodologically flawed. The Turnaway Study addressed the flaws of earlier research by collecting data on baseline mental health conditions and outcomes following either an abortion or a birth when abortion was wanted but denied⁽⁴³⁾. The study followed women for five years, with twice yearly interviews to assess their psychological wellbeing. Being denied an abortion was associated with a greater risk of experiencing adverse psychological outcomes. However, psychological wellbeing improved over time, so that both women who had received and those who had been denied an abortion had similar outcomes⁽⁴³⁾.

The psychological outcome of abortion is optimised when women are able to make decisions based on the complexities of their own intimate partner relationships, and their lives and values ⁽⁴⁴⁾. Women exposed to intimate partner violence are more likely to have an unintended pregnancy ⁽⁴⁵⁾ and those who go on to give birth with an unintended pregnancy are more likely to be the victim of violence from the male involved in the pregnancy than those who have an abortion ⁽⁴⁶⁾.

Medical professionals can assist by being aware that women seeking an abortion might have been exposed to violence or require mental health care for factors that preceded the request for termination ⁽⁴⁷⁾.

See KEMH Women's Health Strategy and Programs [Family and Domestic Violence Toolbox](#) for more information on intimate partner violence and resources.



Risks of carrying a pregnancy to term

Pregnancy and birth are for the majority of healthy women 'low risk' events. There are, however, risks associated with pregnancy, birth and the puerperium.

Mortality risk

Maternal deaths in Australia are rare but healthy women do still die in pregnancy and following birth.

In the decade from 2009 to 2018, there were 251 women reported to have died during pregnancy or within 42 days of the end of pregnancy with a maternal mortality rate of 6.7 deaths per 100 000 women giving birth. Maternal deaths are categorised as either direct or indirect. Direct deaths are the result of obstetric complications or pregnancy or its management, while indirect deaths are the result of conditions without an obstetric cause but that were aggravated by the pregnancy.

The most frequent causes of maternal death in the decade 2009-2018 was pre-existing cardiovascular disease and non-obstetric haemorrhage, then suicide ⁽⁴⁸⁾. The most frequent causes of direct maternal death in the same period were thromboembolism and obstetric haemorrhage.

Morbidity

As the rate of maternal mortality has declined in high-resources countries over the past 50 years, there has been increased emphasis on maternal morbidity. It has been suggested that maternal morbidity is underestimated as there is a tendency to focus on obstetric complications during labour and birth, such as haemorrhage, sepsis, hypertensive disorders and obstructed labour, while other issues such as depression, incontinence, sexual health issues and pelvic girdle pain are underreported ⁽⁴⁹⁾.

In an attempt to construct a shared understanding of maternal morbidity, the World Health Organisation's Maternal Morbidity Working Group has defined maternal morbidity as "any health condition attributed to and/or complicating pregnancy and childbirth that has a negative impact on the woman's wellbeing and/or functioning" ⁽⁵⁰⁾. However, there is currently no standard classification or national database for the collection of morbidity data. Maternal morbidity varies in both duration and severity and covers a wide range of diagnoses ranging from the near death of a woman from complications during pregnancy or childbirth, to non life-threatening illness that impacts on wellbeing more broadly ⁽⁵⁰⁾.

A recent Australian study used the Victorian Perinatal Data Collection (VPDC) database to examine risk factors associated with maternal morbidity and found that lower socioeconomic status, indigenous status, older maternal age and clinical factors such as primiparity, coexisting medical conditions, previous caesarean section and previous pregnancy loss all increase the likelihood of severe maternal morbidity ⁽⁵¹⁾.

Pregnancy and birth data from the Midwives Notification System, which compiles information on all births in Western Australia, recorded pregnancy complications in 30.3 percent of women ⁽⁵²⁾. In 2015, hypertension occurred in 4.1 percent of pregnancies, while the most common complication in those who have given birth were gestational diabetes (8.8 percent) and premature rupture of membranes (3.5 percent) ⁽⁵²⁾. However, longer term consequences following birth that largely remain uncaptured, such as post traumatic-stress disorder, postpartum depression, physical and emotional disabilities and sexual dysfunction, may lead to a significant reduction in quality of life ⁽⁵³⁾.

Other risks associated with pregnancy

Pre-existing problems can be exacerbated by pregnancy. However, only a small proportion of women within the obstetric population have a pre-existing disease.

Women at higher risk of medical and obstetric complications include those with the following factors:

- Obesity
- Diabetes and other endocrine disease
- Cardiovascular disease
- Asthma and other chronic respiratory disease
- Depression
- Other systemic and chronic illnesses
- Smoking, alcohol and other drug consumption in pregnancy
- Previous obstetric complications

Complications in pregnancy and birth range from minor symptoms, such as heartburn, to more serious events, such as major haemorrhage, sepsis, pulmonary embolus, and cardiac failure.

Table 1: Overview of medical risks of abortion and continuing the pregnancy

Risks common to pregnancy and abortion	
Topics	Suggested discussion points
Process	Blood tests, pregnancy tests, ultrasound, costs
Delivery/procedure	Haemorrhage, infection, retained products of conception Drug reactions Injury to the uterus and cervix Retained products
Anaesthetic issues	Method (general anaesthetic, local anaesthetic or twilight sedation) and possible associated risks Conscious (twilight) sedation: Generally associated with less risk than with general anaesthetic but uncommonly respiratory depression can occur and some memory of the event may remain General anaesthesia: Nausea, fever and rare anaesthetic complications Rhesus incompatibility, medical complications
Risks of pregnancy only	
Topics	Suggested discussion points (examples only, not a complete list)
Medical risks of pregnancy such as:	Hypertension, pre-eclampsia, spontaneous miscarriage, antepartum haemorrhage, placenta praevia and rare complications such as a molar pregnancy
Pre-existing systemic diseases	Cardiovascular, respiratory, endocrine, genitourinary and other systemic diseases can place the pregnant woman at greater risk during pregnancy and these diseases can be exacerbated by pregnancy
Fetal conditions	Antibody-incompatibilities, congenital conditions
Physical complications of delivery such as:	Tears to cervix, vagina and perineum due to delivery process and assistance from birthing staff Obstructed labour, Caesarean section and its complications
Problems following delivery	A number of complications can arise, such as: • Depression, PTSD, sexual dysfunction • Infection, of urinary or genital tract or breast • Secondary haemorrhage • Thromboembolic disease Dyspareunia due to scar tissue from tears or episiotomy • Long-term damage to pelvic floor supports with potential for prolapse of uterus, bowel and bladder
Risks of abortion only	
Short term problems	Failure of termination requiring further procedures
Longer term problems	Longer term problems of miscarriage and preterm birth in subsequent pregnancies can occur with multiple induced abortions

Adoption

Information on adoption is provided as some women may wish to consider adoption as an alternative to parenting or abortion.

Adoption practices are shaped by society, culture, religion, politics and economics and have changed over time. Adoption has significantly declined in Australia for a variety of reasons, including increased support for single parent families, the emergence of family planning and legislative changes which provide alternative legal options. There were 310 adoptions in Australia in 2018-19; 57 were adopted from overseas and 253 were adoptions within Australia. Of these, 211 were known child adoptions (ie: adopted by a step parent, relative or carer) and 42 were local adoptions. There has also been a significant shift away from the secrecy that was associated with adoption to a transparent system which focuses on the needs of the child ⁽⁵⁴⁾.

Since 1995, all adoptions in Western Australia have occurred within a policy of 'openness' where the birth mother is involved in each aspect of the adoption. Current research indicates that continued contact increases the birth mother's satisfaction with the process ⁽⁵⁵⁾.

The Turnaway Study, a five-year longitudinal study of 956 women seeking abortion care in the United States of America, including 231 women denied abortion due to gestational limit, found that amongst women seeking abortion, adoption is infrequently chosen ⁽⁵⁶⁾. It has been suggested that women tend to choose adoption when there are fewer options available ⁽⁵⁶⁾. Women should be advised that abortion, parenting, exploration of kin support and adoption are all potential options ⁽⁵⁵⁾.

For more information on adoption see the [Department of Community Development, 'Pregnant and considering adoption for your child?'](#)



Guidelines for medical practitioner counselling

Counselling about the medical risk of termination and pregnancy is required as part of obtaining informed consent. However, counselling for matters relating to the termination of pregnancy and carrying a pregnancy to term must be offered but is not required. The medical practitioner can complete the counselling themselves. Medical practitioners are able to complete non-directive pregnancy counselling training, which will enable them to access Medicare benefits for up to three non-directive pregnancy support counselling services per patient, per pregnancy, from any of the following items - 792, 4001, 81000, 81005 and 81010. Medical practitioners can access non-directive pregnancy counselling training through the Royal Australian College of General Practitioners.

For medical practitioners who look after women considering abortion and provide general health care, it is useful to understand general counselling approaches and principles.

- Every woman who has an unintended and/or unwanted pregnancy requires access to counselling that is confidential and is responsive to her social, emotional and cultural circumstance.
- Counselling must be non-directive and non-judgmental, delivered by professionals who are aware of their own values and attitudes and are ready to refer to another practitioner if there is a conflict which may prejudice the counselling process.
- The purpose of counselling is to assist the woman (and partner where appropriate) to clarify issues surrounding the pregnancy and to come to a decision about the pregnancy outcome and how it is to be achieved.
- Women should be given the opportunity to tell their story, paying attention to their relationships, their support networks and their beliefs about abortion. This process clarifies special needs, vulnerabilities and issues in the decision.
- Options which can be discussed are: a) continuing with the pregnancy, parenting the child alone or with her partner; b) continuing the pregnancy and relinquishing the child for adoption/ fostering; and c) abortion. Include consideration of emotional consequences for all three options.
- In exploring options, help the woman to identify her strengths, her social resources, her belief systems, her needs, issues relating to significant others, and the short and long-term implications of the decision, as well as practical considerations.
- Women who remain ambivalent or undecided should be offered further counselling

Once a decision and plan are made, the woman should be assisted with the implementation and any potential consequences. It is essential that adequate notes are made for clinical and legal purposes.

Resources

Abortion providers

Clinics have different criteria, such as gestation age and medical or surgical abortion

Metropolitan

Burton Street Family Practice
(medical)

23 Burton Street
Bentley WA 6102
Phone (08) 9458 4558

East Fremantle Medical Centre
(medical)

12 Silas Street
East Fremantle WA 6158
Phone (08) 9339 4116

Fiona Stanley Hospital
(medical and surgical)

- » Conditions that would preclude access at community clinics
- » Medical abortion above 63 days

11 Robin Warren Drive
Murdoch WA 6150
Phone (08) 6152 2222

King Edward Memorial Hospital
(medical and surgical)

- » Conditions that would preclude access at community clinics
- » Medical abortion above 63 days
- » Less than 14 years old

Family Services Coordinator at King Edward Memorial Hospital (KEMH)
KEMH.Referrals@health.wa.gov.au
Phone (08) 6458 2222
Fax (08) 64581031

Fremantle Women's Health Centre
114 South Street
Fremantle WA 6160
Phone (08) 9431 0500

Marie Stopes Midland
(medical and surgical)

Free for patients in the St John of God Midland Public Hospital catchment area.
8 Sayer St
Midland WA 6056
Phone 1300 003 707

Nanyara Medical Group
(medical and surgical)

2 Cleaver Terrace
Rivervale WA 6103
Phone (08) 9277 6070

Sexual Health Quarters (SHQ)

70 Roe Street,
Northbridge WA 6003
Phone (08) 9227 6177

Women's Health and Family Services
(medical)

www.whfs.org.au
Phone (08) 6330 5400
1800 998 5400
(free call outside of Perth metro area)

Country WA

Marie Stopes (medical)

Via telehealth 1300 405 568

This service may be an option in many country areas but requires that the patient has access to 24-hour medical care within two hours of home.

Kimberley

Broome Hospital
(medical and surgical)

Robinson Street
Broome WA 6725
Phone (08) 9194 2222

Unplanned pregnancy counselling

Inform if requesting unplanned pregnancy counselling so that the appointment is prioritised. The below services have received funding from the Department of Health to provide non-directive unplanned pregnancy counselling.

Sexual Health Quarters (SHQ)

www.shq.org.au/clinic/unintended-pregnancy/

Phone (08) 9227 6177

Desert Blue Connect (Geraldton)

www.desertblueconnect.org.au/service/unplanned-pregnancy-counselling/

Phone (08) 9964 2742

Goldfields Women's Health Care Centre

(Kalgoorlie)

<https://www.gwhcc.org.au/services/unplanned-pregnancy-counselling/>

Phone (08) 9021 8266

South West Women's Health & Information Centre (Bunbury)

www.swhic.com.au/services/

Freecall 1800 673 350

(08) 9791 3350

Legal services

Legal Aid

Children's Court Protection Service

(for assistance with applications to the Children's Court)

Phone (08) 9218 0160

Youth Legal Service

www.youthlegalserviceinc.com.au

Perth Metro (08) 9202 1688

Pilbara

Karratha Medical Centre

(medical)

5 Sharpe Avenue

Karratha WA 6714

Phone (08) 9185 3555

Mid West

Geraldton Health Campus

(medical and surgical)

51-85 Shenton Street

Geraldton WA 6530

Phone (08) 9956 2222

Victoria District Medical Centre

(medical)

151 Durlacher Street

Geraldton WA 6530

Phone (08) 9921 6099

Goldfields

Kalgoorlie Health Campus

(surgical)

15 Piccadilly Street

Kalgoorlie WA 6433

Phone: (08) 9080 5888

South West

Choices South West

(medical and surgical)

Dunborough Medical Centre

4/54 Dunn Bay Road

Dunborough WA 6281

Phone (08) 9746 3300

Women's Health Services

Fremantle Women's Health Centre

www.fwhc.org.au

Phone (08) 9431 0500

Ishar Multicultural Women's Health Services (Mirrabooka)

www.ishar.org.au

Phone (08) 9345 5335

Midland Women's Health Care Place

www.mwhcp.org.au

Phone (08) 9250 2221

South Coastal Women's Health Services (Rockingham)

Phone (08) 9550 0900

Women's Health and Family Services (Northbridge and Joondalup)

www.whfs.org.au

Phone (08) 6330 5400

1800 998 5400

(freecall outside of Perth metro area)

Women's Health and Wellbeing Services (Gosnells)

www.whfs.org.au

Phone (08) 9490 2258

Desert Blue Connect (Geraldton)

Phone (08) 9964 2742

Kalgoorlie

Goldfields Women's Health Care Centre

www.gwhcc.org.au

Phone (08) 9021 8266

Port Hedland

Hedland Well Women's Centre

www.wellwomens.com.au

Phone (08) 9140 1124

Tom Price

Centre for Women's Safety and Wellbeing

www.cwsw.org.au/services/nintirri-centre

Phone (08) 9189 1556

0456 802 061

Bunbury

South West Women's Health & Information Centre

www.swwhic.com.au

Phone: (08) 9791 3350

Freecall 1800 673 350

Mental Health Services

Beyond Blue

1300 224 636

www.beyondblue.org.au/

[get-support/get-immediate-support](http://www.beyondblue.org.au/get-support/get-immediate-support)

Lifeline

www.lifeline.org.au/

13 11 14

Diverse sexualities and genders

Qlife

www qlife.org.au/get-help

1800 184 527

Living Proud

www.livingproud.org.au/about/

Another Closet

www.anothercloset.com.au/

Other contacts

King Edward Memorial Hospital – Family
services coordinator

KEMH.Referrals@health.wa.gov.au

Phone (08) 6458 2222 (weekdays)

Fax (08) 6458 1031

www.kemh.health.wa.gov.au

Adoption Services

5 Newnan Court

Fremantle WA 6160

Phone (08) 9286 5200

Free Call 1800 182 178

and ask to speak to the local adoptions

duty officer.

[www.wa.gov.au/organisation/departments-](http://www.wa.gov.au/organisation/departments-of-communities/pregnant-and-considering-adoption-your-child)

[of-communities/pregnant-and-considering-](http://www.wa.gov.au/organisation/departments-of-communities/pregnant-and-considering-adoption-your-child)

[adoption-your-child](http://www.wa.gov.au/organisation/departments-of-communities/pregnant-and-considering-adoption-your-child)

Women's Domestic

Violence Helpline

Support and counselling for women

experiencing family and domestic violence,

including referrals to women's refuges.

Phone (08) 9223 1188

1800 007 339

Sexual Assault Resource Centre

Crisis counselling over the phone from

8.30am to 11pm any day of the week.

You can also request a

counselling appointment.

Phone (08) 6458 1828

1800 199 888

Appendix A

Suggested further information points for counselling on the processes involved

Abortion

Topics	Suggested discussion points
Pre-abortion process	Blood tests, pregnancy tests, ultrasound, costs
Abortion	
Anaesthetic issues	Method (GA, LA or twilight) and possible associated risks
Procedure: Type	Medical – mifepristone & misoprostol Surgical – suction/vacuum aspiration
Procedure: General	Waiting period, duration of procedure, recovery time, where performed
Post abortion	
Additional support	Resources on where to access more information about counselling, pregnancy and adoption

Pregnancy

Topics	Suggested discussion points
Pre-delivery	
Pregnancy care process	<ul style="list-style-type: none">• Blood tests, pregnancy tests, ultrasound, costs• Schedule of visits• Antenatal care options• Costs involved
Delivery	
Additional supports	Resources on where to access more information about counselling, pregnancy and adoption
Postnatal	Resources on where to access more information about counselling, postnatal care

Appendix B

Abortion referral process template

Abortion/termination of pregnancy referral	
Patient details	
Patient circumstances	
<p>LMP: _____</p> <p>Pregnancy test results to date (ie: urine, serum, ultrasound) _____</p> <p>Obstetric history:</p> <p>STI screen offered: <input type="checkbox"/> Yes <input type="checkbox"/> No</p> <p>Cervical screening history/offer if due: _____</p>	<p>To ensure the patient is free from coercion, see her alone and discuss:</p> <p><input type="checkbox"/> Circumstances of pregnancy, eg: contraception, assault</p> <p><input type="checkbox"/> Family and community support</p> <p><input type="checkbox"/> Options with unintended pregnancy, ie: continuation of the pregnancy, abortion and adoption</p> <p><input type="checkbox"/> Her relationship with her partner</p> <p>If patient is under 16 years of age seek agreement for her parent/guardian to be involved in the counselling process.</p>
<p>Informed consent for referral to abortion services</p> <p>It is a legal requirement that informed consent for referral to abortion services is obtained by a general practitioner who is not the doctor performing or assisting with the abortion. This is done by:</p> <ul style="list-style-type: none"> • Providing information about the medical risks associated with having an abortion and continuing with the pregnancy • Offering a referral for counselling prior to the abortion and for continuing with the pregnancy (Note: medical practitioners may also provide this counselling however an offer of referral must be made) • Informing the patient that counselling is available post-termination or post-delivery 	
<p>Abortion/termination of pregnancy referral</p> <p>Have you:</p> <p><input type="checkbox"/> Explained both medical and surgical methods of abortion?</p> <p><input type="checkbox"/> Referred to the family services coordinator at KEMH if she has financial concerns in obtaining an abortion? KEMH.referrals@health.wa.gov.au or fax 64581 031, telephone contact via switchboard (08) 6458 2222, and ask for family services coordinator.</p> <p><input type="checkbox"/> Discussed contraception?</p> <p><input type="checkbox"/> Arranged follow-up for seven to 14 days after the abortion?</p>	
<p>Note that clinics have different criteria and different costs. See <u>Health Pathways</u>. Referrals for patients with medical or surgical comorbidities, under 14 years of age and those who are unsuitable for external clinics should be made to King Edward Memorial Hospital (KEMH).</p>	


Notification by medical practitioner of induced abortion

Guide for completing E-form version of Form 1, Notification by medical practitioner of induced abortion

To complete and submit the E-form, go to:

- Link available under "Related Links" at: https://ww2.health.wa.gov.au/en/Articles/N_R/Notification-of-terminations-of-pregnancy-induced-abortion, and
- https://ww2.health.wa.gov.au/Articles/A_E/Abortion-Notification-System.

To seek more information, go to https://ww2.health.wa.gov.au/Articles/A_E/Abortion-Notification-System - this website has contact details, including email address or phone number of maternal and child health team, for assistance if required.




References

1. Moseson H, Herold S, Filippa S, Barr-Walker J, Baum SE, Gerdts C. Self-managed abortion: a systematic scoping review. *Best Practice & Research Clinical Obstetrics & Gynaecology*. 2020;63:87-110.
2. Lee JH, Park HN, Kim NS, Park H-J, Park S, Shin D, et al. Detection of illegal abortion-induced drugs using rapid and simultaneous method for the determination of abortion-induced compounds by LC-MS/MS. *Chromatographia*. 2019;82(9):1365-71.
3. Harris LH, Grossman D. Complications of unsafe and self-managed abortion. *New England Journal of Medicine*. 2020;382(11):1029-40.
4. Gynaecologists TRAaNZCoOa. The use of mifepristone for medical abortion. 2019.
5. Goldstone P, Walker C, Hawtin K. Efficacy and safety of mifepristone-buccal misoprostol for early medical abortion in an Australian clinical setting. *Australian and New Zealand Journal of Obstetrics and Gynaecology*. 2017;57(3):366-71.
6. Kapp N, Lohr PA. Modern methods to induce abortion: Safety, efficacy and choice. *Best Practice & Research Clinical Obstetrics & Gynaecology*. 2020;63:37-44.
7. Mazza D, Burton G, Wilson S, Boulton E, Fairweather J, Black KI. Medical abortion. *Australian Journal of General Practice*. 2020;49(6):324.
8. Gerdts C, Dobkin L, Foster DG, Schwarz EB. Side effects, physical health consequences, and mortality associated with abortion and birth after an unwanted pregnancy. *Women's Health Issues*. 2016;26(1):55-9.
9. Raymond EG, Grimes DA. The comparative safety of legal induced abortion and childbirth in the United States. *Obstetrics & Gynecology*. 2012;119(2):215-9.
10. Excellence NIHaC. Abortion care. 2019.
11. Kerns J, Steinauer J. Management of postabortion hemorrhage. *Contraception*. 2013;87(3):331-42.
12. Costescu D, Guilbert E, Bernardin J, Black A, Dunn S, Fitzsimmons B, et al. Medical abortion. *Journal of Obstetrics and Gynaecology Canada*. 2016;38(4):366-89.
13. Achilles SL, Reeves MF. Prevention of infection after induced abortion. *Contraception*. 2011;83(4):295-309.
14. Excellence NIHaC. Best practice in comprehensive abortion care. 2015.
15. Organization WH. Medical management of abortion. *World Health Organization*; 2019.
16. Gynaecologists RCoO. The care of women requesting induced abortion. 2011.
17. Lichtenberg ES, Paul M. Surgical abortion prior to 7 weeks of gestation. *Contraception*. 2013;88(1):7-17.
18. Gynaecologists TRAaNZCoOa. Guidelines for the use of Rh(D) immunoglobulin (Anti-D) in obstetrics. 2019.
19. Gynaecologists TRAaNZCoOa. COVID-19: Anti D and Abortion 2020 [Available from: <https://ranzcoog.edu.au/news/covid-19-anti-d-and-abortion>].
20. Kapp N, Eckersberger E, Lavelanet A, Rodriguez MI. Medical abortion in the late first trimester: a systematic review. *Contraception*. 2019;99(2):77-86.
21. Freeman MD, Porat N, Rojansky N, Elami-Suzin M, Winograd O, Ben-Meir A. Physical symptoms and emotional responses among women undergoing induced abortion protocols during the second trimester. *International Journal of Gynecology & Obstetrics*. 2016;135(2):154-7.
22. Brouns JFGM, van Wely M, Burger MP, van Wijngaarden WJ. Comparison of two dose regimens of misoprostol for second-trimester pregnancy termination. *Contraception*. 2010;82(3):266-75.
23. Tramer M, Moore A, McQuay H. Propofol anaesthesia and postoperative nausea and vomiting: quantitative systematic review of randomized controlled studies. *British Journal of Anaesthesia*. 1997;78(3):247-55.
24. Kehlet H, Dahl JB. Anaesthesia, surgery, and challenges in postoperative recovery. *The Lancet*. 2003;362(9399):1921-8.
25. Ghoneim M, Mewaldt S. Benzodiazepines and human memory: a review. *Anesthesiology (Philadelphia)*. 1990;72(5):926-38.
26. Wong C, Ng E, Ngai S, Ho P. A randomized, double blind, placebo-controlled study to investigate the use of conscious sedation in conjunction with paracervical block for reducing pain in termination of first trimester pregnancy by suction evacuation. *Human Reproduction*. 2002;17(5):1222-5.
27. Castleman L, Mann C. Manual vacuum aspiration (MVA) for uterine evacuation: Pain management. Chapel Hill, NC, Ipas. 2002.

28. O'Shea LE, Lord J, Fletcher J, Hasler E, Cameron S. Cervical priming before surgical abortion up to 13+ 6 weeks' gestation: A systematic review and meta-analyses for the National Institute of Health and Care Excellence (NICE)—new clinical guidelines for England. *American Journal of Obstetrics & Gynecology MFM*. 2020;100220.
29. Jonathan L, HASLER E, CAMERON S. Cervical priming before surgical abortion between 14 and 24 weeks: A systematic review and meta-analyses for the National Institute for Health and Care Excellence (NICE)—new clinical guidelines for England. *American Journal of Obstetrics & Gynecology MFM*. 2020;100283.
30. Shaw KA, Lerma K. Update on second-trimester surgical abortion. *Current Opinion in Obstetrics and Gynecology*. 2016;28(6):510-6.
31. Carlsson I, Breiding K, Larsson P-G. Complications related to induced abortion: a combined retrospective and longitudinal follow-up study. *BMC women's health*. 2018;18(1):158.
32. Health Nf, Excellence C. Abortion care: National Institute for Health and Care Excellence; 2019.
33. Averbach SH, Seidman D, Steinauer J, Darney P. Re: Prior uterine evacuation of pregnancy as independent risk factor for preterm birth: a systematic review and metaanalysis. *American Journal of Obstetrics & Gynecology*. 2017;216(1):87.
34. Smikle C, Yarrarapu SNS, Khetarpal S. Asherman syndrome. *StatPearls [Internet]*. 2020.
35. Meaidi A, Friedrich SJ, Lidegaard Ø. Risk of surgical evacuation and risk of major surgery following second-trimester medical abortion in Denmark: a nationwide cohort study. *Contraception*. 2020.
36. Obstetricians RCo, Gynaecologists. The care of women requesting induced abortion: RCOG press; 2011.
37. Society AC. Abortion and breast cancer risk 2014 [Available from: <https://www.cancer.org/cancer/cancer-causes/medical-treatments/abortion-and-breast-cancer-risk.html>].
38. Biggs MA, Gould H, Foster DG. Understanding why women seek abortions in the US. *BMC women's health*. 2013;13(1):29.
39. Taft AJ, Powell RL, Watson LF, Lucke JC, Mazza D, McNamee K. Factors associated with induced abortion over time: secondary data analysis of five waves of the Australian Longitudinal Study on Women's Health. *Australian and New Zealand journal of public health*. 2019;43(2):137-42.
40. Rocca CH, Kimport K, Roberts SC, Gould H, Neuhaus J, Foster DG. Decision rightness and emotional responses to abortion in the United States: A longitudinal study. *PloS one*. 2015;10(7):e0128832.
41. Rocca CH, Samari G, Foster DG, Gould H, Kimport K. Emotions and decision rightness over five years following an abortion: An examination of decision difficulty and abortion stigma. *Social Science & Medicine*. 2020;248:112704.
42. Hanschmidt F, Linde K, Hilbert A, Riedel J, Heller SG, Kersting A. Abortion stigma: a systematic review. *Perspectives on sexual and reproductive health*. 2016;48(4):169-77.
43. Biggs MA, Upadhyay UD, McCulloch CE, Foster DG. Women's mental health and well-being 5 years after receiving or being denied an abortion: A prospective, longitudinal cohort study. *JAMA psychiatry*. 2017;74(2):169-78.
44. Chibber KS, Biggs MA, Roberts SC, Foster DG. The role of intimate partners in women's reasons for seeking abortion. *Women's Health Issues*. 2014;24(1):e131-e8.
45. Tinglöf S, Högberg U, Lundell IW, Svanberg AS. Exposure to violence among women with unwanted pregnancies and the association with post-traumatic stress disorder, symptoms of anxiety and depression. *Sexual & Reproductive Healthcare*. 2015;6(2):50-3.
46. Roberts SC, Biggs MA, Chibber KS, Gould H, Rocca CH, Foster DG. Risk of violence from the man involved in the pregnancy after receiving or being denied an abortion. *BMC medicine*. 2014;12(1):144.
47. Horvath S, Schreiber CA. Unintended pregnancy, induced abortion, and mental health. *Current psychiatry reports*. 2017;19(11):77.
48. Welfare AloHa. Maternal death in Australia 2015-2017. 2020.
49. Moran PS, Wuytack F, Turner M, Normand C, Brown S, Begley C, et al. Economic burden of maternal morbidity—A systematic review of cost-of-illness studies. *PloS one*. 2020;15(1):e0227377.
50. Chou D, Tunçalp Ö, Firoz T, Barreix M, Filippi V, von Dadelszen P, et al. Constructing maternal morbidity—towards a standard tool to measure and monitor maternal health beyond mortality. *BMC pregnancy and childbirth*. 2016;16(1):45.

Abortion care: Information and legal obligations for medical practitioners






Government of Western Australia
North Metropolitan Health Service
Women and Newborn Health Service



Abortion:

An information booklet
for consumers



NM



Disclaimer

All information and content in this publication is provided in good faith by North Metropolitan Health Service and is based on sources believed to be reliable and accurate at the time of development. To the fullest extent permitted by law, North Metropolitan Health Service, the Department of Health and the State of Western Australia and its officers, employees and agents are released from liability (including in respect of negligence) for any loss, damage, cost and expense (regardless of whether the loss is direct, indirect or consequential) caused by the use of, or reliance on, this publication and the material contained in it.

This resource is an aid only, and must not be relied upon. This booklet does not replace medical, legal or other

professional advice. Such professional advice should be sought as required and relevant to your particular circumstances. The law is sometimes complex and open to interpretation and applies to different factual circumstances in different ways. If you have a legal problem, you should seek legal advice on the particular circumstances.

Copyright

Copyright to this material is vested in the State of Western Australia unless otherwise indicated. Apart from any fair-dealing for the purpose of private study, research, criticism or review, as permitted under the provisions of the Copyright Act 1968, no part may be reproduced or re-used for any purposes whatsoever without the written permission of the State of Western Australia.

Contents

What is this booklet about?	4
What is abortion?	4
Is abortion safe?.....	5
How do I access an abortion?	5
How much does an abortion cost?.....	6
What happens before an abortion?.....	6
Types of abortion	7
Medical abortion	7
Surgical abortion.....	7
Abortion overview	8
What if I am under 16?	10
What if I am 20 weeks pregnant or more?	10
Counselling and other support	10
Contacts	11

What is this booklet about?

Many women experience an unintended pregnancy. Some will choose to have an abortion. Deciding if you want to have an abortion can be a difficult choice. Every woman has the right to make her own choice about whether to have an abortion or continue with the pregnancy, as long as legal requirements are met.


This booklet provides general information about abortion and encourages you to talk to your health care provider about your options. It is best to approach a health care provider as early in pregnancy as possible. This booklet does not replace medical advice.

Please see the back of this booklet for contact numbers of counselling and support services. Please note that the information in this booklet refers to pregnant women but it is also applicable to other individuals who can become pregnant such as girls and those who are gender diverse.

What is abortion?

An abortion, also known as termination of pregnancy, is a way of ending a pregnancy. This option is available for women in Western Australia who are up to 20 weeks pregnant. After 20 weeks, abortion is only available if the woman or fetus has a severe condition and the procedure must be approved by a panel of medical practitioners, who are specifically appointed for this role.

There are different procedures for abortion. These are covered later in the booklet.



Is abortion safe?

When performed by a qualified health professional, an abortion is usually a safe medical procedure.

Generally, the earlier you have an abortion the safer it is. This is why it is important to discuss your pregnancy and options with a health care provider as early as possible.

Your health care provider will discuss possible complications of having an abortion or continuing the pregnancy. Although serious complications are not common, all medical and surgical procedures have some risks. This will depend on your specific circumstances.

How do I access an abortion?

You will need to make an appointment with your health care provider who will:

- Confirm the pregnancy
- Discuss the medical risks of abortion and pregnancy
- Offer an option of referral to free counselling to assist with decision making, information and support
- Inform you that counselling is also available after an abortion. You can decide if you want to take up the counselling services that are offered
- Refer you to an abortion provider.

Most health care providers will discuss this with you. However, some may not provide consultation and referral on abortion. In this case, you can:

- Approach a different health care provider
- Seek help from one of the Women's Health Services (see list on page 13) or
- Contact Sexual Health Quarters on (08) 9227 6177 for information and referral options.


How much does an abortion cost?


The cost of the abortion will depend on how far along you are in the pregnancy, whether you have a medical or surgical abortion and where you have the procedure. Some costs are covered through Medicare and the fees will be less if you are eligible for Medicare benefits. Talk to your health care provider if you are unable to afford an abortion as you might be able to receive financial assistance via King Edward Memorial Hospital (KEMH).

What happens before an abortion?

Before an abortion, you will need to give your informed consent for referral to an abortion provider. You will then be referred for an abortion. Your health care provider will ask questions about your medical history and may do some tests and screening, which can include blood tests, screening for sexually transmitted infections and an ultrasound to confirm how many weeks pregnant you are.

You will be given information on the types of abortion available to you, what to do after the procedure and pain relief options, as well as information on relevant support services





Types of abortion

There are two types of abortion: medical (where medicines are used) and surgical (where an operation is performed). Most women will generally be able to choose between these; however, this will also depend on how many weeks pregnant you are and your individual circumstances.

Medical abortion

Medical abortion uses a combination of two medications to end a pregnancy. It is available up to 63 days of pregnancy via a general practitioner or community abortion provider. After 63 days it is only available in a hospital setting.

The first tablet (mifepristone) is usually taken when you are at the clinic with your health care provider. It blocks a pregnancy hormone. To complete the abortion, you will be given a second tablet (misoprostol) to take 36 to 48 hours later. It causes the uterus (womb) lining to break down and you will start to bleed, usually within two hours. The pregnancy ends with bleeding similar to a miscarriage. This usually happens within four to six hours of taking the second tablet but may be quicker or take longer.

If, for any reason, this treatment does not work, you may require a surgical termination and further costs may be involved. The abortion service will provide you with information and contact details so you can access help if there are any complications.

Surgical abortion


A surgical termination usually involves a vacuum aspiration and will be done at a clinic or hospital. The procedure is performed under anaesthetic. Occasionally women do not have sedation and instead a local anaesthetic may be used to numb the cervix (neck of the womb). For women over 13 weeks, a slightly different process is used to ensure all the products of pregnancy are removed.

Abortion overview

The table below provides a summary of medical and surgical abortion. You should talk to your doctor or health care provider before deciding which one is right for you.

Medical abortion

What happens in this type of abortion?	This procedure uses medication, given in doses over two or more days that induce miscarriage-like symptoms.
Where will the abortion take place?	It can take place in a clinic or at home.
How long does it take?	You pass the contents of pregnancy within four to six hours of taking the second medicine. Bleeding may take several days to settle.
Is the abortion painful?	You may have some pain (mild to severe) and bleeding or clotting (like a heavy period). Pain relief medication, as ordered by your health care provider, should help with any discomfort.
How much bleeding will there be?	You will have vaginal bleeding during a medical abortion. This is usually heavier than your normal period. After passing the pregnancy, bleeding can last for about 12 days but should get lighter day by day.



Surgical abortion

What happens in this type of abortion?

This is usually performed by an operation called a suction curette under anaesthetic as a day procedure. In more advanced pregnancy (after 13 weeks), the procedure is technically different, called dilation and evacuation.

Where will the abortion take place?

It will take place in a clinic or hospital as a day procedure.

How long does it take?

It usually takes less than 20 minutes, depending on how far along your pregnancy is. However, you should expect to spend up to several hours for the whole process if your care is in a clinic. If your care is in a hospital, you may need to stay for several hours or overnight on a ward.

Is the abortion painful?

Some cramping or mild abdominal pain after the procedure is normal and pain relief medication (ordered by your health care provider) or a hot pack should help with any discomfort.

How much bleeding will there be?

You will have vaginal bleeding after surgical abortion. Bleeding can last for 10-14 days but should get lighter day by day.

What if I am under 16?

If you are under 16 and still financially dependent on a parent/legal guardian, then your parent/legal guardian must be given the opportunity to participate in counselling and consultation about the proposed abortion with you and your health care provider.

If you do not want your parent/legal guardian to know about your pregnancy, you can apply for an order at the Children's Court. Legal assistance for this is free. The legal service will help you with the application and putting your case to the magistrate, who will make the decision on whether to involve your parent/guardian. The Children's Court Protection Service, Legal Aid can be contacted on (08) 9746 3300.

What if I am 20 weeks pregnant or more?

If your pregnancy is 20 weeks or more, abortion is unavailable except in specific circumstances, such as you or the fetus having a severe medical condition. Your health care provider can discuss this with you. They can also request advice from King Edward Memorial Hospital to assist with making an application on your behalf for abortion at 20 weeks and over to the Ministerial Termination Panel, which oversees this process.

Counselling and other support

The decision to have an abortion is yours alone. It can be helpful to talk to someone supportive and unbiased. The unplanned pregnancy counselling services listed in this booklet offer counselling to assist with decision making and post-abortion counselling. These services are free.

Research shows that the majority of women cope well after an abortion. Some women may find the experience stressful or difficult. However, for most women there are no long term psychological or mental health impacts.

Domestic and sexual abuse

There are some circumstances where seeking an abortion might be linked to other distressing events, such as pregnancy through unwanted sexual contact and domestic violence.

You can discuss these issues with your health care provider or women's health service.

Contacts

Abortion providers

(Clinics have different criteria, eg: pregnancy weeks and medical or surgical.)

Metropolitan

King Edward

Memorial Hospital

(Medical and surgical)

Provides abortion services to women who are unable to access private abortion services due to medical risk factors, and women seeking medical abortion after nine weeks gestation.

274 Bagot Road Subiaco WA 6008

Phone (08) 9340 2222

Marie Stopes Midland

(Medical and surgical)

Free for patients in the St John of God Midland Public Hospital catchment.

8 Sayer Street

Midland WA 6056

Phone 1300 003 707

Women's Health and Family Services

(Medical)

www.whfs.org.au

Phone (08) 6330 5400

1800 998 399

(freecall outside Perth metro area)

Sexual Health Quarters (SHQ)

70 Roe Street

Northbridge WA 6003

Phone (08) 9227 6177

www.shq.org.au

Burton Street Family Practice

(Medical)

23 Burton Street

Bentley WA 6102

Phone (08) 9458 4558

East Fremantle Medical Centre

(Medical)

12 Silas Street

East Fremantle WA 6158

Phone (08) 9339 4116

Fiona Stanley Hospital

(Medical and surgical)

11 Robin Warren Drive,

Murdoch WA 6150

Nanyara Medical Group

(Medical and surgical)

2 Cleaver Terrace

Rivervale WA 6103

Phone (08) 9277 6070

Country WA

Marie Stopes

(Medical)

Via telehealth 1300 405 568

This service may be an option in country areas but requires that the patient has access to 24-hour medical care within two hours of home.

Kimberley

Broome Hospital

(Medical and surgical)

Robinson Street Broome WA 6725

Phone (08) 9194 2222

Pilbara

Karratha Medical Centre

(Medical)

5 Sharpe Avenue

Karratha WA 6714

Phone (08) 9185 3555

Mid West

Geraldton Health Campus

(Medical and surgical)

51 Shenton Street Geraldton WA 6530

Phone (08) 9956 2222

Victoria District Medical Centre

(Medical)

151 Durlacher Street

Geraldton WA 6530

Phone (08) 9921 6099

Goldfields

Kalgoorlie Health Campus

(Surgical)

15 Piccadilly Street

Kalgoorlie WA 6433

Phone (08) 9080 5888

South West

Choices South West

(Medical and surgical)

Dunsborough Medical Centre

4/54 Dunn Bay Road

Dunsborough WA 6281

Phone (08) 9746 3300

Legal services

Legal Aid

Children's Court Protection Service

(for assistance with applications to the Children's Court)

Phone (08) 9218 0160

Youth Legal Service

www.youthlegalserviceinc.com.au

Perth Metro

Phone (08) 9202 1688

Unplanned pregnancy counselling

Please advise the service if you are requesting unplanned pregnancy counselling so that your appointment is prioritised, or if you need an appointment for counselling following an abortion.

Sexual Health Quarters (SHQ)

70 Roe Street,
Northbridge WA 6003
Phone (08) 9227 6177
www.shq.org.au

Desert Blue Connect (Geraldton)

www.desertblueconnect.org.au/
Phone (08) 9964 2742

Goldfields Women's Health Care Centre (Kalgoorlie)

[www.gwhcc.org.au/services/
unplanned-pregnancy-counselling/](http://www.gwhcc.org.au/services/unplanned-pregnancy-counselling/)
Phone (08) 9021 8266

South West Women's Health & Information Centre (Bunbury)

Freecall 1800 673 350
Phone (08) 9791 3350
www.swwhic.com.au/services/

Other support services

Women's Domestic Violence Helpline

Support and counselling for women experiencing family and domestic violence, including referrals to women's refuges.

Phone (08) 9223 1188
1800 007 339

Sexual Assault Resource Centre

Crisis counselling over the phone from 8.30am to 11pm any day of the week.

You can also request a counselling appointment

Phone (08) 6458 1828
1800 199 888

Women's Health Services

Fremantle Women's Health Centre

www.fwhc.org.au
Phone (08) 9431 0500

Ishar Multicultural Women's Health Services (Mirrabooka)

www.ishar.org.au
Phone (08) 9345 5335

Midland Women's Health Care Place

www.mwhcp.org.au
Phone (08) 9250 2221

South Coastal Women's Health Services (Rockingham)

Phone (08) 9550 0900

Women's Health and Family Services (Northbridge and Joondalup)

www.whfs.org.au
Phone (08) 6330 5400
1800 998 5400 (freecall outside Perth metro area)

Women's Health and Wellbeing Services (Gosnells)

www.whws.org.au
Phone (08) 9490 2258

Desert Blue Connect (Geraldton)

www.desertblueconnect.org.au
Phone (08) 9964 2742



Goldfields Women's Health Care Centre (Kalgoorlie)

www.gwhcc.org.au

Phone (08) 9021 8266

Hedland Well Women's Centre

www.wellwomens.com.au

Phone (08) 9140 1124

Centre For Women's Safety and Wellbeing (Tom Price)

www.csws.org.au/services/nintirri-centre

Phone (08) 9189 1556

0456 802 061

South West Women's Health & Information Centre (Bunbury)

www.swwhic.com.au

Phone: 08 9791 3350

Freecall 1800 9791 3350

Mental Health Services

Beyond Blue

Phone 1300 22 4636

www.beyondblue.org.au/get-support/get-immediate-support

Lifeline

www.lifeline.org.au/

Phone 13 11 14

Diverse sexualities and genders

Qlife

www qlife.org.au/get-help

Phone 1800 184 527

Living Proud

www.livingproud.org.au/about

Another Closet

www.ssdv.acon.org.au




Sexual Health Quarters (SHQ)

70 Roe Street,


Northbridge WA 6003

www.shq.org.au

Phone (08) 9227 6177







Women and Newborn Health Service

King Edward Memorial Hospital

374 Bagot Road, Subiaco WA 6008

☎ (08) 6458 2222

🌐 wnhs.health.wa.gov.au

This document can be made available
in alternative formats on request.

© North Metropolitan Health Service 2022

